

## **Recommendations for Volunteer Potato Control:**

Growers should minimize the number of tubers left in the field during potato harvest (Steiner et al., EB1993). Newberry and Thornton (2004) demonstrated that deep fall tillage (mold board plow) that buries tubers deeper prior to cold winter temperatures should be avoided. Previous studies comparing tillage practices indicated that plowing **following** a deep penetrating frost could be beneficial by exposing deeper buried tubers to additional freezing events (Thomas and Smith, 1983). However, weather patterns are not always conducive to make this practice effective and field access in winter months is often limited.

Control measures should strive to minimize competition with rotational crops and formation of new daughter tubers that can persist and cause problems in subsequent crops. Several components of volunteer potato management that growers can implement in this year's rotational crops are listed below.

- On higher value crops with nematode problems such as carrot or onions, spring fumigate with metham sodium (Vapam, Busan, and others) and 1, 3,-dichloropropene (Telone II). Field studies indicate about 70 to 75% of tubers are killed by a combination of Telone II at 10 GPA applied with shanks plus Vapam at 30 GPA applied by center pivot. Lower rates of fumigants are less effective in killing tubers. Follow labels for proper rates, soil temperatures, soil moisture, and time required between fumigation and planting of subsequent crop.
- If possible, delay planting of the rotation crop to allow maximum early volunteer potato emergence and apply glyphosate (Roundup) or remove with tillage. If planting glyphosate resistant crops, wait to apply glyphosate until the majority of the volunteer potatoes have emerged and are just beginning to initiate tubers.
- Maleic hydrazide applied to the potato crop can significantly reduce volunteer potato sprouting the following year. Check with processor before using.
- Use herbicides that are active in reducing volunteer potatoes in rotation crops. Several herbicides can be very effective in killing potato plants and reducing daughter tuber weight, including mesotrione (Callisto), fluroxypyr (Starane), atrazine (Aatrex, Atrazine), glyphosate (Roundup), dicamba + diflufenzopyr (Distinct), dicamba (Banvel, Clarity), and imazamox (Raptor). Ethofumesate (Nortron) applied preemergence retards potato emergence and applied postemergence can reduce potato competition with the crop. Repeated applications of contact herbicides such as, oxyfluorfen (Goal), carfentrazone (Aim), fomesafen (Reflex), glufosinate (Rely), and paraquat (Gramoxone) can also be effective. Follow labels closely for labeled crops, proper rates, timing of applications, and crop rotation restrictions.
- When possible, apply postemergence herbicides when potatoes are just beginning to initiate tubers on stolons. If applications are made earlier, mother tubers often resprout and the volunteer plants will require additional herbicide applications. If applications are made later, yield loss may have

already occurred and many new tubers will have already formed which will infest next year's crop.

- Previous USDA-ARS research demonstrated that cultivation about 1 week after postemergence applications of Starane, Goal, Roundup, and Banvel greatly reduced the number of daughter tubers formed compared to herbicides alone. In corn, Callisto herbicide has reduced new daughter tuber formation greater than other postemergence herbicides. Cultivation after Callisto application may not improve volunteer potato control.
- Select competitive crops and those with effective herbicide and cultivation options like field corn. Crops like carrots have few effective herbicides registered for volunteer potato control, so avoid planting such crops in fields where volunteers will be plentiful. Winter wheat is a very competitive crop and delays volunteer potato emergence in the spring. However, cultivation isn't practical in wheat and there are limited opportunities for timing effective herbicide applications in winter wheat prior to new tuber set on volunteer potatoes.
- Repeated cultivations and hand weeding can control volunteer potatoes, but they are most effective and economical when combined with other control methods.
- Grazing fields with hogs, sheep, or cattle may also reduce the number of tubers available to sprout.

For more detailed control information contact Rick Boydston, USDA-ARS, 24106 N Bunn Road, Prosser, WA 99350. Phone (509) 786-9267. Email: [rick.boydston@ars.ars.usda.gov](mailto:rick.boydston@ars.ars.usda.gov)

#### References.

1. Boydston, R. A. Volunteer potato (*Solanum tuberosum*) control with herbicides and cultivation in field corn (*Zea mays*). Weed Tech. 15:461-466. 2001.
2. Boydston, R. A. Managing volunteer potato (*Solanum tuberosum*) in field corn (*Zea mays*) with carfentrazone-ethyl and dicamba. Weed Technol. 18:83-87. 2004.
3. Boydston, R. A., H. P. Collins, and A. K. Alva. Control of volunteer potato (*Solanum tuberosum*) in sweet corn with mesotrione unaffected by atrazine and tillage. Weed Technol. 22:654-659. 2008.
4. Boydston, R. A. and M. D. Seymour. Volunteer potato control (*Solanum tuberosum*) with herbicides and cultivation in onion (*Allium cepa*). Weed Technol. 16:620-626. 2002.
5. Boydston, R. A. and M.M. Williams II. Effect of fumigation on volunteer potato (*Solanum tuberosum*) tuber viability. Weed Technol. 17:352-357. 2003.
6. Boydston, R. A. and M. M. Williams, II. Managing volunteer potato (*Solanum tuberosum*) in field corn with mesotrione and arthropod herbivory. Weed Technol. 19:443-450. 2005.
7. Newberry, GD and RE Thornton. 2004. Influence of post harvest tillage and rotation crop selection on volunteer potato survival. Abstract in Amer. Potato J

81:77.

8. Steiner, C. M., G. Newberry, R. Boydston, J. Yenish, and R. Thornton. Volunteer potato management in the Pacific Northwest rotational crops. Washington State University Extension Publications, EB1993, 12 pp. 2005.
9. Thomas, PE and DR Smith. 1983. Relationship between cultural practices and the occurrence of volunteer potatoes in the Columbia Basin in northwest United States. Amer Potato J 60:289-294.
10. Williams, M. M. II and R. A. Boydston. Alternatives to handweeding volunteer potato (*Solanum tuberosum*) in carrot (*Daucus carota*). Weed Technol. 19:1050-1055. 2005.
11. Williams II, M. M. and R. A. Boydston. Effect of shoot removal during tuberization on volunteer potato (*Solanum tuberosum*) tuber production. Weed Technol. 16:617-619. 2002. Williams, M. M. II and R. A. Boydston. Volunteer potato (*Solanum tuberosum* L.) interference in carrot. Weed Science 54:94-99. 2006.